



Brief report

Body Mass Index, Overweight, and Obesity in Swedish Women Born Post-term

José G. B. Derraik,^a Maria Lundgren,^b Wayne S. Cutfield,^a Fredrik Ahlsson^b

^aLiggins Institute, University of Auckland, Auckland, New Zealand

^bDepartment of Women's and Children's Health, Uppsala University, Uppsala, Sweden

Abstract

Background: There is increasing evidence that post-term birth (≥ 42 weeks of gestation) is associated with adverse long-term outcomes. We assessed whether women born post-term displayed increased risk of overweight and obesity in adulthood.

Methods: Data were collected at first antenatal visit (~10–12 weeks of gestation) on singleton Swedish women aged ≥ 18 years in 1991–2009 (mean age 26.1 years), who were born post-term ($n = 27\ 153$) or at term (37–41 weeks of gestation; $n = 184\ 245$). Study outcomes were evaluated for continuous associations with gestational age. Stratified analyses were carried out comparing women born post-term or at term. Analyses were also run with a 2-week buffer between groups to account for possible errors in gestational age estimation, comparing women born very post-term (≥ 43 weeks of gestation; $n = 5761$) to those born within a narrower term window (38–40 weeks of gestation; $n = 130\ 110$).

Results: Increasing gestational age was associated with greater adult weight and body mass index (BMI). Stratified analyses showed that women born post-term were 0.5 kg heavier and had BMI 0.2 kg/m² greater than those born at term. Differences were more marked between women born very post-term (≥ 43 weeks) vs. a narrower term group (38–40 weeks): 1.0 kg and 0.3 kg/m². The adjusted relative risks of overweight/obesity and obesity in women born very post-term were 1.13 and 1.12 times higher, respectively, than in those born at term.

Conclusions: Post-term birth is associated with greater BMI and increased risk of overweight and obesity in adulthood, particularly among women born ≥ 43 weeks of gestation.

Keywords: adult, body mass index, females, gestational age, obesity, overweight, prolonged gestation.

Marked physiological stress during pregnancy is commonly associated with a pattern of adverse metabolic sequelae in the offspring, which can be detected in childhood. This is principally manifested as adverse effects on insulin sensitivity, which has been shown to be reduced in children born too small,¹ too large (from diabetic mothers),² or too early (preterm).³

Recently, it was observed that prepubertal children born post-term (≥ 42 weeks of gestation) had a 34% reduction in insulin sensitivity, as well as increased body fat and abdominal adiposity.⁴ Further, longitudinal data from Sweden showed that males born post-term had accelerated weight gain during childhood and an associated increased risk of obesity in adoles-

cence.⁵ Thus, it was of interest to examine whether adverse outcomes seen in childhood and adolescence persist into adulthood. We aimed to assess whether women born post-term displayed increased risk of overweight and obesity in a large cohort of mothers with data from early pregnancy.

Methods

In Sweden, information is collected prospectively during pregnancy from the first antenatal visit and is subsequently forwarded to the Swedish Birth Register, which was established in 1973 and contains data on >99% of births in the country.⁶ This study was approved by the Regional Ethical Review Board in Uppsala.

We examined data from the first antenatal visit (mostly 10–12 weeks of gestation) on all singleton

Correspondence:

José G. B. Derraik, Liggins Institute, University of Auckland, Private Bag 92019, Auckland, New Zealand.

E-mail: j.derraik@auckland.ac.nz

Swedish women aged ≥ 18 years in 1991–2009. Exclusion criteria were birth to non-Nordic mothers, congenital malformations, or preterm birth (< 37 weeks of gestation). Further, women were only included if also born to a singleton mother aged ≥ 18 years at their first antenatal visit. For women with two or more pregnancies in the study period, data were only included for the first recorded pregnancy.

Gestational age (extracted from the Swedish Birth Register) was mostly based on the maternal report of the last menstrual period, otherwise determined from ultrasound scans. We compared women born post-term (≥ 42 weeks of gestation) vs. those born at term (37–41 weeks of gestation). However, due to potential inaccuracies associated with gestational age estimation in the second trimester,⁷ it is very likely that a number of women born at term would have been misclassified as post-term. As a result, we also examined differences in the two groups with a 2-week buffer between them, comparing women born very post-term (≥ 43 weeks of gestation) to those born within a narrower term window (38–40 weeks of gestation).

Weight was measured and current height was self-reported or measured, with body mass index (BMI) being subsequently calculated. Overweight/obesity was defined as BMI ≥ 25.0 kg/m² and obesity as ≥ 30.0 kg/m².

Study outcomes (i.e. anthropometry) were evaluated for continuous associations with gestational age, while stratified analyses were also carried out comparing the above-described groups. Analyses were performed using general linear regression models, accounting for a number of confounding factors: sibship size (i.e. total number of siblings), birth order, age, year of birth, SGA status (SGA by birthweight and/or length vs. not SGA), maternal smoking during pregnancy, and family situation (two-parent family vs. other arrangements). Generalised linear models with a similar construction were run to evaluate binary outcomes (e.g. obesity risk). Statistical analyses were carried out in SAS v.9.4 (SAS Institute, Cary, NC, USA). Where applicable, results are expressed as adjusted relative risks (aRR) or β coefficients with associated 95% confidence intervals.

Results

There were 303 301 Swedish women who were born within 1973–1988 and gave birth to a child within

1991–2009. A total of 65 298 women failed to meet the inclusion criteria and a further 26 605 women had incomplete anthropometric data in early pregnancy. Thus, we examined data on 211 398 women at an average of 26.1 years of age (range 18–36 years), including 27 153 women born post-term and 184 245 women born at term.

Continuous associations

Increasing gestational age was associated with greater adult weight [in kg, $\beta = 0.127$ (0.087–0.166)] and BMI [$\beta = 0.030$ (0.016–0.043)] in women. Weight and BMI were lowest in women born at 39 weeks of gestation, progressively increasing with greater gestational age (Figure 1).

Stratified analyses

Women born post-term were 0.5 kg heavier and had a BMI that was 0.2 kg/m² greater than those born at term (Table 1). Women born post-term also had a slightly greater risk of being overweight/obese (aRR 1.05) or obese (aRR 1.08) (Table 1).

However, when a 2-week buffer was applied between groups, differences were more marked. Women born very post-term (≥ 43 weeks of gestation) were 1.0 kg heavier and had a BMI 0.3 kg/m² greater than those born within a narrower term window (38–40 weeks) (Table 2). The adjusted relative risk of




Figure 1. Body mass index (BMI) early in pregnancy (10–12 weeks of gestation) among 211 398 Swedish women, according to their gestational age at birth. Data are means and 95% confidence intervals, adjusted for confounding factors (i.e. sibship size, birth order, age, year of birth, family situation, SGA status, and maternal smoking during pregnancy).

Table 1. Anthropometric data recorded early in pregnancy (10–12 weeks of gestation) among Swedish women born post-term (≥ 42 weeks of gestation) or at term (37–41 weeks of gestation)

	Unadjusted		Adjusted		P-value
	Term	Post-term	Term	Post-term	
<i>n</i>	184 245	27 153	184 245	27 153	
Age (years)	26.1 \pm 3.9	26.3 \pm 3.9			
Height (cm)	167.2 \pm 6.0	167.3 \pm 5.9	164.7 (164.6, 164.9)	164.9 (164.7, 165.1)	0.21
Weight (kg)	67.4 \pm 12.8	67.8 \pm 12.9	65.3 (64.9, 65.7)	65.8 (65.4, 66.2)	<0.0001
BMI (kg/m ²)	24.1 \pm 4.3	24.2 \pm 4.3	24.0 (23.9, 24.2)	24.2 (24.0, 24.3)	<0.0001
Overweight/obese (BMI ≥ 25.0 kg/m ²)	31.5%	32.8%			
	1.00	1.04 (1.02, 1.06)	1.00	1.05 (1.03, 1.07)	<0.0001
Obese (BMI ≥ 30.0 kg/m ²)	9.5%	10.0%			
	1.00	1.05 (1.01, 1.09)	1.00	1.08 (1.04, 1.12)	<0.001

Unadjusted data are percentages, means \pm standard deviations, or relative risks and 95% confidence intervals. Adjusted data are means or relative risks with respective 95% confidence intervals, adjusted for confounding factors (i.e. sibship size, birth order, age, year of birth, family situation, SGA status, and maternal smoking during pregnancy).

overweight/obesity was 1.13 and of obesity 1.12 times higher in women born very post-term (Table 2).

Discussion

This study shows that women born post-term were heavier, had greater BMI, and had increased risk of overweight and obesity early in pregnancy. Notably, there was a progressive increase in BMI with increasing gestational age from a nadir at 39 weeks of gestation, so that the risk of overweight and obesity was particularly higher in those born very post-term (≥ 43 weeks of gestation).

Despite changes in obstetric practice and lack of data for many countries, there is still a great variation in the rates of post-term births across the world. It has been estimated that post-term births account for as few as 0.4% of live births in Austria to as many as 8.1% in Denmark⁸ and 5.5% in the USA.⁹ Thus, individuals born post-term are a relatively common group in some countries. However, post-term birth had not been identified as a risk factor for adverse metabolic outcomes until very recently.^{4,5} Apart from these studies in childhood and adolescence,^{4,5} it seems that the possible effects of post-term birth on anthropometry and metabolism in adulthood have not been

Table 2. Anthropometric data recorded early in pregnancy (10–12 weeks of gestation) among Swedish women born very post-term (≥ 43 weeks of gestation) or at a narrower term window (38–40 weeks of gestation)

	Unadjusted		Adjusted		P-value
	Term	Very post-term	Term	Very post-term	
<i>n</i>	130 110	5761	130 110	5761	
Age (years)	26.0 \pm 3.9	26.5 \pm 4.0			
Height (cm)	167.2 \pm 6.0	167.2 \pm 5.9	164.7 (164.5, 164.9)	164.8 (164.6, 165.1)	0.07
Weight (kg)	67.3 \pm 12.7	68.0 \pm 13.1	65.2 (64.7, 65.6)	66.2 (65.6, 66.7)	<0.0001
BMI (kg/m ²)	24.1 \pm 4.3	24.3 \pm 4.3	24.0 (23.8, 24.1)	24.3 (24.1, 24.5)	<0.0001
Overweight/obese (BMI ≥ 25.0 kg/m ²)	31.3%	34.6%			
	1.00	1.11 (1.07, 1.15)	1.00	1.13 (1.09, 1.17)	<0.0001
Obese (BMI ≥ 30.0 kg/m ²)	9.4%	10.1%			
	1.00	1.07 (0.99, 1.16)	1.00	1.12 (1.03, 1.21)	0.006

Unadjusted data are percentages, means \pm standard deviations, or relative risks and 95% confidence intervals. Adjusted data are means or relative risks with respective 95% confidence intervals, adjusted for confounding factors (i.e. sibship size, birth order, age, year of birth, family situation, SGA status, and maternal smoking during pregnancy).

examined.¹⁰ Thus, our findings are of interest, as the observed increased risk of overweight and obesity in adult women born post-term may be also associated with greater cardio-metabolic risk. Although the proportion of infants born at 43 weeks and beyond seems to be much reduced (maybe 0.1–1.5% of all births in Europe⁸ and 21.2% of all post-term births in our study), our findings suggest that this group may be particularly at risk of adverse outcomes.

Notably, sexual dimorphism has been commonly observed in studies examining the effects of early life programming on long-term health.¹¹ Mathai *et al.*¹² showed in a small New Zealand study that men born preterm were heavier and had greater BMI than term counterparts in their mid-30s, while these parameters were similar amongst women. For those born at the other end of the gestational age spectrum, Beltrand *et al.* observed that the adverse effects of post-term birth were markedly greater among male adolescents.⁵ As a result, the magnitude of the associations observed in women born post-term cannot be automatically extrapolated to men due to possible sexual dimorphism, and it would be important to evaluate whether there are more marked associations among men born post-term.

The factors underpinning the metabolic programming in children and adults born post-term are unclear, but may be associated with genetic inheritance or an adverse fetal environment in late gestation.⁴ It is possible that post-term infants may be exposed to physiological stress due to an abnormally long gestation or late pregnancy utero-placental insufficiency.¹⁰ Placentae from post-term pregnancies have abnormal histological features that are similar to placentae of babies born small-for-gestational-age, including apoptotic changes, diffuse calcifications, chorionic villus degeneration, infarcts, increased necrosis, and decreased perfusion surface.^{10,13,14} However, undernutrition *in utero* is unlikely to occur in most prolonged pregnancies, since post-term and term infants have comparable ponderal indices and adjusted birth weights.⁴ Thus, adverse outcomes may not be necessarily associated with the extended period *in utero per se*.¹⁰ There may be underlying genetic or epigenetic mechanisms determining both the prolonged gestation and the associated metabolic programming.¹⁰

The limitations of our study include the lack of data on socioeconomic status and other potential confounding factors, particularly maternal BMI. In

Sweden, the odds of post-term delivery among obese women are 1.63 times higher than in normal-weight women.¹⁵ Therefore, one cannot discard the possibility that maternal overweight/obesity may be an important factor underpinning increased BMI among women born post-term. In addition, there are inaccuracies associated with gestational age estimation using the date of last menstrual period, which are less precise than early ultrasound scans and overestimate the number of post-term births.⁷ The misclassification of women born at term as post-term could have masked stronger associations between post-term birth and risks of overweight and obesity in adulthood. This also highlights the importance of our stratified analyses including a 2-week buffer between groups to account for possible errors in gestational age estimation. Another limitation was our relatively homogenous population of Nordic women, so that findings may not be readily extrapolated to other ethnic groups.

In conclusion, post-term birth was associated with greater BMI and increased odds of overweight and obesity in adult women. However, the relevance of these findings for later cardiovascular and metabolic risks is unclear. Further, other studies need to identify whether or not maternal overweight/obesity is an important driver of the observed associations. Future studies should also assess whether the pattern and magnitude of such associations differ in adult men born post-term.

Competing interest

The authors have no competing financial or non-financial interests to declare.

Source of funding

There has been no specific funding received by the authors to support this study.

References

- Hofman PL, Cutfield WS, Robinson EM, Bergman RN, Menon RK, Gluckman PD. Insulin resistance in short children with intrauterine growth retardation. *Journal of Clinical Endocrinology and Metabolism* 1997; 82:402–406.
- Hunter WA, Cundy T, Rabone D, Hofman PL, Harris M, Regan F, *et al.* Insulin sensitivity in the offspring of women with type 1 and type 2 diabetes. *Diabetes Care* 2004; 27:1148–1152.

- 3 Hofman PL, Regan F, Jackson WE, Jefferies C, Knight DB, Robinson EM, *et al.* Premature birth and later insulin resistance. *New England Journal of Medicine* 2004; 351:2179–2186.
- 4 Ayyavoo A, Derraik JGB, Hofman PL, Mathai S, Biggs J, Stone P, *et al.* Pre-pubertal children born post-term have reduced insulin sensitivity and other markers of the metabolic syndrome. *PLoS ONE* 2013; 8:e67966.
- 5 Beltrand J, Soboleva TK, Shorten PR, Derraik JGB, Hofman P, Albertsson-Wikland K, *et al.* Post-term birth is associated with greater risk of obesity in adolescent males. *Journal of Pediatrics* 2012; 160:769–773.
- 6 Cnattingius S, Ericson A, Gunnarskog J, Kallen B. A quality study of a medical birth registry. *Scandinavian Journal of Social Medicine* 1990; 18:143–148.
- 7 Savitz DA, Terry JW, Dole N, Thorp JM, Siega-Riz AM, Herring AH. Comparison of pregnancy dating by last menstrual period, ultrasound scanning, and their combination. *American Journal of Obstetrics and Gynecology* 2002; 187:1660–1666.
- 8 Zeitlin J, Blondel B, Alexander S, Breart G. Variation in rates of postterm birth in Europe: reality or artefact? *BJOG* 2007; 114:1097–1103.
- 9 Martin JA, Hamilton BE, Osterman MJ, Curtin SC, Matthews TJ. Births: final data for 2013. *National Vital Statistics Reports* 2015; 64:1–65.
- 10 Ayyavoo A, Derraik JGB, Hofman PL, Cutfield WS. Postterm births: are prolonged pregnancies too long? *Journal of Pediatrics* 2014; 164:647–651.
- 11 Gabory A, Roseboom TJ, Moore T, Moore LG, Junien C. Placental contribution to the origins of sexual dimorphism in health and diseases: sex chromosomes and epigenetics. *Biology of Sex Differences* 2013; 4:5.
- 12 Mathai S, Derraik JGB, Cutfield WS, Dalziel SR, Harding JE, Biggs J, *et al.* Increased adiposity in adults born preterm and their children. *PLoS ONE* 2013; 8:e81840.
- 13 Mihu CM, Mihu D. Placental, morphological, and ultrasonographical particularities in post-term pregnancies. *Leonardo Electronic Journal of Practices and Technologies* 2004; 4:51–61.
- 14 Smith SC, Baker PN. Placental apoptosis is increased in post-term pregnancies. *BJOG* 1999; 106:861–862.
- 15 Roos N, Sahlin L, Ekman-Ordeberg G, Kieler H, Stephansson O. Maternal risk factors for postterm pregnancy and cesarean delivery following labor induction. *Acta Obstetrica et Gynecologica Scandinavica* 2010; 89:1003–1010.